


Chronic Diseases: a global view

Surveillance/NMH



WORLD

Disease Burden (DALYs), by broad cause group, 2001



Source: WHO 2002



Leading Causes of Mortality and Burden of Disease

Preliminary estimates for 2000

Mortality

%

- Ischaemic heart disease 13.7
- Cerebrovascular disease 9.5
- Lower respiratory infections 6.4
- HIV/AIDS 4.2
- COPD 4.2
- Diarrhoeal diseases 4.1
- Perinatal conditions 4.0
- Tuberculosis 2.8
- Lung Cancer 2.3
- Road traffic accidents 2.2

DALYs


%

- Lower respiratory infections 6.7
- HIV/AIDS 6.2
- Perinatal conditions 6.2
- Diarrhoeal diseases 5.0
- Depression 4.1
- Ischaemic heart disease 4.1
- Cerebrovascular disease 3.5
- Malaria 3.1
- Road traffic accidents 2.8
- COPD 2.7



CHINA : Leading Causes of Death in Rural Areas, 1998

Rural Population: 813 million



 Communicable diseases (2.6%)


 Noncommunicable conditions (82.9%)

 Injuries (11.2%)


 Undiagnosed (3.3%)




Trends in CHD deaths in urban and rural China (age standardized)




Ischaemic heart disease




Stroke




Ischaemic heart disease



Stroke



Cardiovascular (CVD) epidemic in countries of different stages of development




Source: WHO, NMH/MNC



Heart Disease and Stroke

More than 12 million deaths each year

 = 100,000 deaths




Source: WHR 2002



Heart Disease and Stroke

 = 100,000 deaths



More than 50% of these deaths can be prevented by reducing major risk factors such as:

- High blood pressure
- High cholesterol
- Low fruit and vegetable intake
- Tobacco use
- Obesity
- Physical inactivity

Source: WHR 2002




2002 World Health Report

“Reducing risks, promoting healthy life”

www.who.int/whr

www.thelancet.com



What is a risk?

**“A probability of an adverse outcome
or a factor that raises this probability”**




Criteria for choosing risk factors

- Likely to be among the leading causes of disease burden
- Not too specific or too broad
- High likelihood of causality
- Reasonably complete data
- Potentially modifiable




Continuous exposure and disease associations


Blood pressure




Cholesterol




Body mass index



Distribution of attributable burden by exposure levels



Distribution of attributable burden by exposure levels



Deaths attributable to leading risk factors, 2000

	Male (000)	Female (000)	Total (000)	%
Blood Pressure	3,491	3,649	7,141	12.8
Tobacco Use	3,893	1,014	4,907	8.8
Cholesterol	2,112	2,303	4,415	7.9
Underweight	1,900	1,848	3,748	6.7
Unsafe Sex	1,370	1,516	2,866	5.1
Low Fruit & Veg	1,449	1,277	2,726	4.9
Overweight	1,168	1,423	2,591	4.6
Physical Inactivity	961	961	1,922	3.4
Alcohol Consumption	1,638	166	1,804	3.2
Unsafe Water/Sanitation	895	835	1,730	3.1



DALYs attributable to leading risk factors, 2000

	Male (000)	Female (000)	Total (000)	%
Underweight	69,733	68,067	137,801	9.6
Unsafe Sex	42,600	49,269	91,869	6.4
Blood Pressure	34,920	29,350	64,270	4.5
Tobacco Use	48,177	10,904	59,081	4.1
Alcohol Consumption	49,397	8,926	58,323	4.0
Unsafe Water/Sanitation	27,432	26,726	54,158	3.8
Cholesterol	22,136	18,301	40,437	2.8
Overweight	15,543	17,872	33,415	2.3
Low Fruit & Veg	15,117	11,544	26,662	1.9
Physical Inactivity	10,159	8,933	19,092	1.3



WHR 2002: Immersed in a sea of risk

Leading 10 selected risk factors as causes of disease burden

■ = Major NCD risk factor

Developing countries		Developed countries
High Mortality	Low Mortality	
1 Underweight	Alcohol	Tobacco
2 Unsafe sex	Underweight	Blood pressure
3 Unsafe water	Blood pressure	Alcohol
4 Indoor smoke	Tobacco	Cholesterol
5 Zinc deficiency	Body mass index	Body mass index
6 Iron deficiency	Cholesterol	Low fruit & vegetable intake
7 Vitamin A deficiency	Iron deficiency	Physical inactivity
8 Blood pressure	Low fruit & vegetable intake	Illicit drugs
9 Tobacco	Indoor smoke from solid fuels	Underweight
10 Cholesterol	Unsafe water	Iron deficiency



Main message: NCD Risk Factors contribute to untimely deaths in all country settings

<i>Condition</i>	<i>Cardiovascular Disease*</i>	<i>Diabetes</i>	<i>Cancer</i>	<i>Chronic- obstructive pulmonary Disease</i>
<i>Risk factor</i>				
<i>Smoking</i>				
<i>Alcohol</i>				
<i>Physical Inactivity</i>				
<i>Nutrition</i>				
<i>Obesity</i>				
<i>Raised Blood pressure</i>				
<i>Dietary fat/Blood lipids</i>				
<i>Blood glucose</i>				

* Including heart disease, stroke, hypertension



Summary

- Small number of risks cause a huge number of chronic disease deaths
- “**Risk transition**” is now occurring in many parts of the world---countries facing a ‘double burden’ of risks i.e. maternal conditions and communicable disease plus noncommunicable diseases
- Substantial decrease in burden of disease possible even with just moderate risk factor modification



WHO's response :

An integrated approach to NCD surveillance and prevention

- **Growing burden of NCD, MH and Injury**
- **Impact of globalisation, urbanisation**
- **Greater emphasis on prevention**
- **Need for standard, comparable data**
- **Greater focus on trends in major risk factors**
- **WHO framework for surveillance (STEPS)**



A framework for Surveillance of major risk factors: a STEPwise approach

- **Hierarchical framework**
- **Standard methods, definitions and protocols**
- **Common approach with core, expanded and optional items**
- **Adaptable to local settings and existing systems**
- **Guiding principle: Keep It Simple (KISS)!**




The WHO STEPwise approach: a framework for Surveillance of major NCD risk factors

Different levels of assessment

- Behaviours
- Physical measurements
- Blood samples

Three modules per risk factor:

- core
- expanded core and
- optional.



WHO STEPS - a tool for surveillance of major NCD risk factors

Step 1: Behaviors


- Tobacco Use
- Physical Inactivity
- Intake fruit/veg
- Alcohol Use

Step 2: Physical measures


- Height /Weight
- Blood Pressure

Step 3: Blood samples


- Blood glucose/diabetes
- Cholesterol



Capacity Building in Developing Countries through Regional Networks



Identification of risk factor data



Identifying useful sources

CountryName

Region Name: WESTERN PACIFIC REGION Source Code: 100288
 100589
 100591
 800341
 800342

Country Name: Japan

Survey

Source_Code	Survey Code	Title	Year start:	Geographic
▶ 100591	100591a1	Kokumin Eiyou Chosa Kekka No Gaiyou	2000	Japan

Update


RF Data

	Sex_N	StartA	EndA	SampleSize	Survey_Code	Best_Record	Graph_Trend	Graph_ASR	Verified
▶	female	20	29	407	100591a1	☒	☐	☐	☒
	female	30	39	589	100591a1	☒	☐	☐	☒
	female	40	49	653	100591a1	☒	☐	☐	☒
	female	50	59	844	100591a1	☒	☐	☐	☒
	female	60	69	725	100591a1	☒	☐	☐	☒
	female	70	100	667	100591a1	☒	☐	☐	☒
	female	20	100	3885	100591a1	☒	☐	☐	☒
	male	20	29	338	100591a1	☒	☐	☐	☒
	male	30	39	394	100591a1	☒	☐	☐	☒
	male	40	49	464	100591a1	☒	☐	☐	☒
	male	50	59	597	100591a1	☒	☐	☐	☒
	male	60	69	655	100591a1	☒	☐	☐	☒
	male	70	100	489	100591a1	☒	☐	☐	☒

Record: 1 of 233



Generating reports



WHO SuRF Report

WHO SuRF Report

Please select a region and a country

Region Name:

Country Name:

View Report

Example



WHO SuRF Report

WHO SuRF Report

Please select a region and a country

Region Name:

Country Name:

View Report *Print Report*



Japan

WESTERN PACIFIC REGION

General Information

	Males	Females
2000 Total Population	62,376,060	65,160,880
2020 Projected Population	60,896,580	65,059,020
2000 Average Life Expectancy (years)	77.5	84.7
Uncertainty interval	77.4 - 77.7	84.4 - 85.1

Risk Factors

Tobacco use


Sex	Age Groups	n	Prevalence	95% CI
Current daily smoker				
male	20 - 29	337	60.8	
	30 - 39	394	56.6	
	40 - 49	463	55.1	
	50 - 59	597	54.1	
	60 - 69	656	37.0	
	= 70	490	29.4	
	= 20	2937	47.4	45.6-49.2
female	20 - 29	407	20.9	
	30 - 39	590	18.8	
	40 - 49	655	13.6	
	50 - 59	844	10.4	
	60 - 69	725	6.6	
	= 70	668	4.0	
	= 20	3889	11.5	10.5-12.5
Survey Year	2000 - 2000			
Survey Population	National; Japan			
Source:	Ministry of Health LaWJ; Kokumin Eiyō Chōsa Kōkai No Gaiyō Results of National Nutrition Study 2001 Nov 8 ;			



WHO Global NCD Infobase

February 12, 2003





Definition: Current daily smoker

Survey Population: National; Japan

Source: Kokumin Eiyō Chōsa Keikaku No Gaiyō/Results of National Nutrition Study, 2001 Nov 8



WHO Global NCD Infobase

February 12, 2003



Conclusions

- Role of established risk factors greater than commonly thought
- In many world regions, the leading 5 risk factors account for more than one-third of mortality and one-quarter of DALYs
- Risks are widespread – all risk factors have global impact, and the burden of many occurs predominantly in developing countries



Conclusions

- Cause(s) known of more than two-thirds of many major diseases eg. ischaemic heart disease, stroke, diabetes
- Research agenda driven by need to identify and implement affordable, practical interventions
- Substantial and rapid improvements in overall healthy life expectancy are possible if:
 - major risks are targeted
 - population-wide changes are achieved

